



# Shoulder Arthroscopy with Distal Clavicle Excision Rehabilitation Protocol

---

Stephanie A. Boden, MD

**PT Frequency:** 2-4 times per week

**ROM Restrictions:** No cross-body/horizontal adduction until 8 weeks

**Brace:** Sling for comfort only 0-2 weeks

## Phase I: Week 0-4

- Sling x 0-2 weeks for comfort only
- 0-2 weeks: Elbow, wrist, hand ROM, shoulder pendulums, wall stretch at home
- 2-4 weeks: Grip strengthening, pulleys/canes, elbow/wrist/hand ROM, pendulums
- Deltoid cuff isometrics
- Begin scapular protraction/retraction
- Passive to Active shoulder ROM as tolerated
- **No** cross-body adduction until 8 weeks

## Phase II: Weeks 4-8

- Increase ROM to full as tolerated
- Advance isometric with arm at side, rotator cuff and deltoid
- Advance to TheraBand and dumbbells as tolerated
- Capsular stretching at end-ROM to maintain flexibility
- **No** cross-body adduction until 8 weeks

## Phase III: Weeks 8-12

- Advance to full, pain-free ROM
- Advance strengthening as tolerated
- Begin eccentrically resisted motion and closed chain activities
- Advance to sport and full activity as tolerated after 12 weeks